



WARNING
HARMFUL · ECOTOXIC
KEEP OUT OF REACH OF CHILDREN



ELITO[®]

FUNGICIDE

A systemic triazole (DMI) fungicide for the control of diseases in cereals, grass seed crops, peas and onions, for the control of brown rot in stonefruit and to control rust and fusarium in pasture.

ACTIVE INGREDIENT: Contains 430 g/litre tebuconazole in the form of a suspension concentrate.

NET CONTENTS: 1 LITRES



HSNO CLASS: 6.1E, 6.9B, 9.1A, 9.3C.**TOXICITY:**

HARMFUL: May be harmful if swallowed, inhaled or absorbed through the skin.
HARMFUL: May cause organ damage (endocrine system) from repeated oral exposure at high doses.

ECOTOXICITY:

VERY TOXIC: Very toxic to aquatic organisms with long lasting effects. Avoid contamination of any water supply with product or empty container.
HARMFUL: Harmful to terrestrial vertebrates.

PRECAUTIONS:**KEEP OUT OF REACH OF CHILDREN.**

When handling, wear eye protection, waterproof gloves, boots and overalls.
 Remove protective clothing and wash hands and face thoroughly before meals and after work.
 Wash protective clothing after use.
 Do not eat, drink or smoke while using.

STORAGE:

Store in the original container tightly closed away from foodstuffs.
 Avoid storage at temperatures below 5°C and above 35°C.
 Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.

DISPOSAL:

Triple rinse empty container and add rinsate to the spray tank.
 Recycle empty container. Otherwise crush and bury empty container in a suitable landfill. Do NOT use the container for any other purpose.
 Avoid contamination of any water supply with chemical or empty container.

SPILLAGE:

Wear appropriate protective clothing and prevent material from entering waterways.
 Contain the spill by damming, recover spill product by absorbing with sawdust or an inert absorbent material then transfer the recovered spill material to a properly labelled container.
 Dispose of waste safely in an approved landfill or other approved facility.

FIRST AID: If swallowed do NOT induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Begin artificial respiration if the victim is not breathing. Use mouth to nose rather than mouth to mouth. **For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.**

TRANSPORT: Do not carry more than 10 litres of this product on a passenger service vehicle.

RECORD KEEPING: Records of use as described in NZS 8409 Management of Agrichemicals must be kept if 3 or more litres of ELITO is used within 24 hours.

MIXING: Shake the container well before use. For large containers roll and invert several times to ensure uniform re-suspension of the contents. Add two thirds of the required quantity of water to the spray tank. With the agitation system running, add the required amount of ELITO to the spray tank. Agitate for 3-5 minutes while adding the remaining water. Continue to agitate while spraying or restarting after a stoppage.

CLEANING: Flush equipment thoroughly with clean water after use. If the spray unit which has applied ELITO is to be used for the application of other chemicals to sensitive crops, ensure the unit is completely decontaminated.

COMPATIBILITY: ELITO is compatible with most commonly used insecticides and fungicides.

RESISTANCE MANAGEMENT:

ELITO contains a fungicide from the triazole group of fungicide and has the DMI (Demethylation Inhibitors) mode of action. Resistance to this fungicide could develop from excessive use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies that exist for any fungus listed on the label. Do not use another product containing a fungicide in the same group in conjunction with ELITO. For further information see: www.resistance.nzmps.org.

DIRECTIONS FOR USE:

It is an offence to use this product on animals.

For field crops ensure complete coverage by applying at least 200 litres water/ha by ground, or 50 litres water/ha by air. Apply preventatively when disease levels are low but disease risk is high. For stonefruit the rates suggested are for high volume spraying to run off. For concentrate spraying adjust dilution rate accordingly.

Rainfall within one hour following application may reduce effectiveness. If 15mm or more rain falls within 3 hours of treatment, pasture should be re-sprayed.

CROP	DISEASE	RATE	NOTES
Barley	Ear disease complex, leaf rust, net blotch (spot and net form), Ramularia, scald, spot blotch	440ml/ha	Apply at first appearance of disease and repeat 3-4 weeks later if disease pressure persists. For the ear disease complex, apply a maximum of 3 DMI applications, including seed treatment, per crop. Where powdery mildew is expected to be severe, use DMI's in conjunction with alternative fungicides.
Wheat	Glume blotch, leaf rust, powdery mildew, speckled leaf blotch, stripe rust, ear disease complex.		
Oats	Crown rust		
Onions	White rot	875ml/ha	Apply in 1000 litres of water/ha. Make the first application 6-8 weeks after sowing and follow with two further applications at 3-4 week intervals.
Peas	Powdery mildew	145ml/ha	Apply preventatively at flowering or at first signs of disease. Repeat if necessary 14 days later.
Ryegrass seed crops	Crown rust, stem rust	440ml/ha	Apply at first appearance of disease and repeat 3-4 weeks later if necessary.
	Blind seed disease	440ml/ha	Apply at the start of flowering and repeat 10-14 days later.
Stonefruit	Brown rot (blossom blight)	23ml/ 100 litres of water. (apply a minimum of 465ml/ha)	Blossom blight: Apply at 7-10 day intervals from pink to shuck fall. Shorter intervals may be needed in wet weather. Apply non DMI fungicides in the pre-harvest period. Brown rot pre-harvest: Apply 2-3 applications at 7-10 day intervals in the pre-harvest period. Use non DMI fungicides earlier in the season. Do not apply more than 3 applications per season, preferably in a mixture with a non DMI fungicide.
Pasture	Rust, fusarium	440ml/ha	Apply at first signs of disease. Ensure complete coverage of grazed areas. A second application may be necessary if conditions favour disease development. Do not apply to the same area at less than a 28 day interval, or more than twice per season. Do not apply when temperatures are above 20oC or wind speeds exceed 12 kph. Ground application: Apply in 100-200 litres of water/ha. Add an organosilicone surfactant at 100ml/100 litres of water. Aerial application: Apply in 50 litres of water/ha. Add an organosilicone surfactant at 200ml/100 litres of water.

WITHHOLDING PERIODS:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

Cereals for grain - 49 days. Fodder cereals - 28 days. Onions - 35 days.

Peas - 14 days. Ryegrass seed crops - 30 days.

Stonefruit - 1 day. For export crops check with Summerfruit NZ or your exporter.

Pasture - Animals (including lactating animals) may be reintroduced into treated pasture 10 days after application. Animals for slaughter must be placed on clean pasture for 6 days prior to slaughter. Pasture is considered clean from residues 45 days after treatment. Where pasture cannot be guaranteed clean, such as in aerial applications, animals must not be sent for slaughter if 51 days after treatment of the pasture.

CONDITIONS OF SALE: The use of ELITO being beyond the control of the suppliers, no Warranty expressed or implied is given by Global Agrichem Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Global Agrichem Ltd accept no responsibility whatsoever resulting from the use of this product.

Registered pursuant to the ACVM Act 1997: P9175. See www.foodsafety.govt.nz for registration conditions. Approved under the HSNO Act 1996: HSR101009. See www.epa.govt.nz for controls.

REGISTERED PROPRIETOR:

Suite 4, level 1, 38 Whitaker place,
Grafton, Auckland, New Zealand.

Tel: +64 9 9729837

FAX: +64 9 3531736

info@agcro.com.nz



GLOBAL AGRICHEM

Date of Manufacture:

Batch No:

ELITO® is a registered trademark of Global Agrichem Ltd.

