



**GLOBAL AGRICHEM**

**WARNING**  
**KEEP OUT OF REACH OF CHILDREN**  
**ECOTOXIC, COMBUSTIBLE LIQUID**



# **GLOBAL GLUFOSINATE<sup>®</sup>**

## **HERBICIDE**

A non-selective herbicide active against a wide range of grasses, broad-leaved weeds and clovers.

**ACTIVE INGREDIENT:** Contains 200 g/l glufosinate -ammonium in the form of a soluble concentrate.

**NET CONTENTS: 20 LITRES**



WORLD CLASS: 6.1E, 6.3B, 6.4A, 6.8B, 6.9A, 9.1C, 9.2B, 9.3C.

## TOXICITY:

TOXIC: May cause organ damage from repeated oral exposure at high doses.  
 HARMFUL: May be harmful if swallowed, inhaled or absorbed through the skin.  
 HARMFUL: May cause reproductive/development damage from repeated oral exposure.  
 HARMFUL: May cause skin irritation.  
 HARMFUL: May cause eye irritation.

## ECOTOXICITY:

HARMFUL: Harmful to aquatic organisms with long lasting effects. Avoid contamination of any water supply with product or empty container.  
 TOXIC: Toxic to the soil environment.  
 HARMFUL: Harmful to terrestrial vertebrates.  
 NON-SELECTIVE HERBICIDE: Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Do not allow drift to occur outside the target area.

## APPROVED HANDLER:

This product must be used under the control of an approved handler when it is applied in a wide dispersive manner or used by a commercial contractor.  
 This product must not be used for any purpose or in any manner contrary to this label unless authorised under appropriate legislation.

## PRECAUTIONS:

### KEEP OUT OF REACH OF CHILDREN.

Keep away from sources of ignition.  
 Avoid contact with skin, eyes and inhalation of spray mist.  
 When handling, wear eye protection, waterproof gloves, boots and overalls.  
 Remove protective clothing and wash hands and face thoroughly before meals and after work.  
 Do not eat, drink or smoke while using.

## STORAGE:

Store in the original container away from sources of ignition in a locked, dry, cool area away from foodstuffs, fertilisers and seeds. Avoid storage at temperatures below 5°C and above 35°C.

Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.

## DISPOSAL:

Triple rinse empty container and add rinseate to the spray tank.  
 Recycle empty container. Otherwise crush and bury empty container in suitable landfill. Do NOT use the container for any other purpose.  
 Avoid contamination of any water supply with chemical or empty container.

## SPILLAGE:

Wear appropriate protective clothing and prevent material from entering waterways.  
 Contain the spill by damming, recover spill product by absorbing with sawdust or an inert absorbent material then transfer the recovered spill material to a properly labelled container. Dispose of waste safely in a suitable landfill.

WITHHOLDING PERIOD: 180 days on all crops.

DIRECTION FOR USE: RATES OF APPLICATION

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

USE	DOSAGE RATE	REMARKS
Berryfruit-Raspberries and Brambles Primocane control	500-750ml/100 litres applied through cone nozzles in sufficient water to ensure complete coverage	Apply as a directed high volume spray to primocanes not exceeding 25cm in length. Avoid drift onto foliage of fruiting canes. Use the high rate on vigorously growing crops. The addition of a non-ionic wetter at label rates is recommended. Knockdown takes approximately 10 days. Do not apply to plants before the first primocanes for the season have reached a height of 15 cm. Do not apply to plants under stress.
Orchard and vineyard weed control- Pipfruit, kiwifruit, stonefruit, grapes and citrus.	Use a dosage rate suitable for the weeds present. If the dilution rate is less than 1 litre /100 litres of water, add a non-ionic wetter at label rates.	Apply in spring when the majority of the annual weeds have germinated. Mix with residual herbicides for long term weed control. Avoid contact with the foliage and immature bark of desirable plants.
Stale seedbed preparation particularly where clover is a problem weed.	3-5 litres/ha in 200-300 litres of water.	Apply to seedling weeds and clover in sufficient water for optimum coverage. May be applied up to one day before crop emergence.
Non-crop land. General weed control.	7.5-10 litres/ha in 300-500 litres of water or 100ml /10 litre knapsack.	Apply as a dranching spray to the point of run-off. Mix with residual herbicides for long term weed control.

## WEED SUSCEPTIBILITY

The lower rates recommended are appropriate for young succulent actively growing weeds or when the weed population is sparse and good coverage and penetration is possible. Use the higher dosage rates for mature weeds or where the weed population is dense.  
 Note: GLOBAL GLUFOSINATE will not completely control mature weeds with well-developed tap roots or underground storage organs. Some regrowth may occur following treatment.

## Dosage Range for major weeds:

3-5 litres/ha			
Annual poa	Clover seedlings	Rough-stalked meadow grass	Subterranean clover
Asparagus seedling	Corbind	Scotch thistle	Wild turnip
Beaked hawkbeard	Daisy	Soft thistle	Willow weed
Black nightshade	Dandelion	Staggergrass	Wireweed
Calandrina	Mouse-ear chickweed	Summer grass	
Chickweed	Musty storksbill	Spurry	
Cleavers	Redroot	Stinking mayweed	
5-7.5 litres/ha			
Bristle grass	Clovers	Parsley piert	Storksbill
Bur medic	Cut-leaved geranium	Prairie grass	Strawberry
Catsear	Docks	Ryegrasses*	Twin cress
Chewings fescue	Fleabane	Sheeps sorrel	Yorkshire sheep
7.5-10 litres/ha			
Broom	Crowfoot grass	Mallows	Wild strawberry
Buttercup's	Gorse	Paspalum	Willow herb
Couch	Hawkbeard	Tall fescue	Yarrow
Field pansy	Kikuyu grass		
10 litres plus/ha			
Bracken	Plantains	Rushes	Indian doab
Meadow rice grass	Rape	Sedges	

\*Resistant weeds: Hairly buttercup and some ryegrass strains.

CONDITIONS OF SALE: The use of GLOBAL GLUFOSINATE being beyond the control of the suppliers, no Warranty expressed or implied is given by Global Agrichem Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Global Agrichem Ltd accept no responsibility whatsoever resulting from the use of this product.

Registered pursuant to the ACVM Act 1997, P9084. See [www.foodsaftey.govt.nz](http://www.foodsaftey.govt.nz) for registration conditions. Approved under the HSNO Act 1996, HSR007813. See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.

REGISTERED PROPRIETOR:



GLOBAL AGRICHEM

Suite 4, level 1, 38 Whitaker place,  
 Grafton, Auckland, New Zealand.  
 Tel: +64 09 9729837  
 FAX: +64 09 3531736  
 info@agro.co.nz



Batch No:

Date of Manufacture:

Global Glufosinate® is a registered trademark of Global Agrichem Ltd.



AGRECOVERY