



**GLOBAL AGRICHEM**

**WARNING**  
**HARMFUL · TOXIC**  
**KEEP OUT OF REACH OF CHILDREN**



# **IMIDATAR<sup>®</sup>**

## **INSECTICIDE**

A systemic insecticide for the control of thrips in onions, mealybug in grapes and aphids in lettuce and vegetable brassica seedlings.

**ACTIVE INGREDIENT:** Contains 600g/litre imidacloprid in the form of a suspension concentrate.

**NET CONTENTS: 1 LITRES**



**HSNO CLASS: 6.1C, 8.3A, 6.5B, 6.9B, 9.1A, 9.2A, 9.3A, 9.4A.**

### TOXICITY:

**DANGER:** May cause eye damage.

**TOXIC:** May be fatal if swallowed, inhaled or absorbed through the skin.

**HARMFUL:** May cause organ damage from repeated oral exposure at high doses.

**HARMFUL:** May cause sensitisation from prolonged skin contact.

### ECOTOXICITY:

**VERY TOXIC:**

- Very toxic to fish with long lasting effects. Avoid contamination of any water supply with product or empty container.

- Very toxic to terrestrial invertebrates.

- Very toxic to the soil environment.

- Very toxic to terrestrial vertebrates.

### PRECAUTIONS:

**KEEP OUT OF REACH OF CHILDREN.**

Avoid contamination of any water supply with product or empty container.

Wash splashes of concentrate from eyes or skin immediately.

When handling or applying, wear full protective clothing gloves and goggles.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Wash protective clothing after use.

Do not eat, drink or smoke while using.

### STORAGE:

Store in the original container tightly closed under lock and key away from feed, seed or foodstuffs.

Do not store with Class 1, 2, 3, 4 or 5 substances.

Stores containing more than 100kg of this product require secondary containment and are subject to signage.

Avoid storage at temperatures below 5°C and above 35°C.

Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.

### DISPOSAL:

Dispose of this product only by using according to this label.

Triple rinse empty container and add rinsate to the spray tank.

Recycle empty container. Otherwise crush and bury empty container in a suitable landfill. Do NOT use the container for any other purpose.

Avoid contamination of any water supply with chemical or empty container.

### SPILLAGE:

Wear appropriate protective clothing and prevent material from entering waterways.

Contain the spill by damming, recover spill product by absorbing with sawdust or an inert absorbent material then transfer the recovered spill material to a properly labelled container.

Dispose of waste safely in an approved landfill or other approved facility.

### RESISTANCE MANAGEMENT:

IMIDATAR contains the Group 4 Insecticide Imidacloprid (Neonicotinoid). Resistance to this insecticide could develop from excessive use. To minimize this risk use strictly in accordance with the label instructions and resistance management strategies that exist for any insect species listed on this label. Do not use another product containing an insecticide in the same group in conjunction with IMIDATAR. For further information see [www.resistance.nzpps.org](http://www.resistance.nzpps.org).

**MIXING:** Shake or roll the container well before use. Add the required quantity of IMIDATAR to a half filled spray tank, then complete filling while agitating. Maintain agitation throughout filling and application.

**COMPATIBILITY:** IMIDATAR is compatible with most commonly used fungicides.

### DIRECTIONS FOR USE:

**It is an offence to use this product on animals.**

CROP	PEST	DOSAGE RATE	REMARKS
Onions	Thrips	130ml in 500-600 litres of water /ha.	Apply when monitoring of thrips levels indicate the threshold level has been reached. Repeat applications at 7-10 day intervals with a maximum of 4 applications. The addition of a wetting agent is suggested. Ensure good spray coverage.
Vegetable brassicas (cell transplants)	Cabbage aphid	5ml in 2.5 litres of water per 1000 plants	Apply evenly in a watering can to seedling trays 1 day before transplanting.
Lettuce (cell transplants)	Lettuce aphid	12-17 ml in 2,5 litres of water per 1000 plants	Apply evenly in a watering can to seedling trays 1 day before transplanting.
Grapes (young non-bearing vines or vines for removal)	Mealybug	0.875ml / vine in 500ml water	Apply as a single drench treatment in autumn or winter to the base of non-bearing vines or vines awaiting removal. If rain or overhead irrigation is not likely in the two days after drenching, apply 2 litres of water per vine once the drench has soaked in. Treated vines must be clearly identified. Any crop set on treated vines must be destroyed within 10 weeks of fruit set.

### WITHOLDING PERIOD:

**It is an offence for the users of this product to cause residues exceeding the relevant MRL in the NZ (Maximum Residues of Agricultural Compounds) Food Standard.**

Onions: 7 days

Vegetable brassicas (cell transplants): 70 days

Lettuce (cell transplants): 35 days

Grapes: Fruit from treated vines MUST be destroyed. No fruit should be harvested for human or animal consumption until the following season.

**CONDITIONS OF SALE:** The use of IMIDATAR being beyond the control of the suppliers, no warranty expressed or implied is given by Global Agrichem Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Global Agrichem Ltd accept no responsibility whatsoever resulting from the use of this product.

Registered pursuant to the ACVM Act 1997, P9177. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions. Approved under the HSNO Act 1996, HSR101008. See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.

**REGISTERED PROPRIETOR:**

Suite 4, level 1, 38 Whitaker place,  
Grafton, Auckland, New Zealand.

Tel: +64 9 9729837

FAX: +64 9 3531736

info@agrcr.co.nz



**GLOBAL AGRICHEM**

Batch No:

Date of Manufacture:

IMIDATAR® is a registered trademark of Global Agrichem Ltd.

